

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004-2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG):

R2SBWYwt

Wyoming Big Sagebrush Semi Desert with Trees

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Gary Back gback@srk.com

Reviewers

Stanley G. Kitchen skitchen@fs.fed.us
Peter Weisberg pweisberg@cabnr.unr.edu

Vegetation Type

Shrubland

Dominant Species*

ARTRW8

CHVI8

ACHY

HECO26

General Model Sources

- Literature
 Local Data
 Expert Estimate

LANDFIRE Mapping Zones

12 17
13 18
16

Rapid Assessment Model Zones

- California Pacific Northwest
 Great Basin South Central
 Great Lakes Southeast
 Northeast S. Appalachians
 Northern Plains Southwest
 N-Cent.Rockies

Geographic Range

This PNVG is found in the southern portion of the Great Basin; western CA, central NV, and UT

Biophysical Site Description

This widespread PNVG is common to the Basin and Range province. In elevation it ranges from 4,500 - 7,000 ft, and occurs on well-drained soils on foothills, terraces, slopes and plateaus. It is found on soil depths greater than 18 inches and up to 60+ inches. Elevationally it is found between low elevation salt desert shrub and mountain big sagebrush zones where pinyon and juniper can establish. Occurs from 4 to 12 inch precipitation zones.

Vegetation Description

Shrub canopy cover generally ranges from 5 to 25%, but can exceed 30% at the upper elevation and precipitation zones. Wyoming big sagebrush sites have fewer understory species relative to other big sagebrush types. Rabbit rubberbrush co-dominant. Perennial forb cover is usually <10%. Perennial grass cover may reach 20 - 25% on the more productive sites. Bluebunch wheatgrass may be a dominant species following replacement fires and as a co-dominant after 20 years. Bottlebrush squirreltail and Indian ricegrass are common. Percent cover and species richness of understory are determined by site limitations. Pinyon (generally *Pinus monophyla*) and juniper (generally *Juniper osteosperma*) present, occasionally reaching 90% canopy cover in areas that have escaped fire. Wyoming big sagebrush semi-desert is critical habitat for the Greater Sage Grouse and many sagebrush obligates.

Disturbance Description

This PNVG is characterized by replacement fires where shrub canopy exceeds 25% (50 - 100 years; mean FRI of 125 years, i.e., 80% of total fire probability) or where grass cover is >15% and shrub cover is > 20% (40 - 70 years; mean FRI of 100 years). Mixed Severity fires account for 20% of fire activity (mean FRI of

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

500 years) where shrub cover ranges from 10 to 20% (20 - 40 years). Surface fires where shrub cover is <10% (0 - 20 years) and generally uncommon during early development (FRI of 200 years). Where pinyon or juniper has encroached after 100 years without fire, mean FRI of fire replacement increases from 100 to 125 years.

The Aroga moth is capable of defoliating large acreages (i.e., > 1,000 ac), but usually 10 to 100 acres.

Weather stress: Prolonged drought (1 in 100 years) on the more xeric sites may reduce shrub cover. Flooding may also cause mortality if the soil remains saturated for an extended period of time (i.e., 1 in 300 year flood events).

Herbivory (non-insect); Herbivory can remove the fine fuels that support Mixed Severity fires and result in woody fuel build up that leads to severe Replacement fires. Surface fires occur in the early seral stage where shrub cover is < 10%.

Adjacency or Identification Concerns

This community may be adjacent to mountain big sagebrush at elevations above 6,500 ft., or adjacent to pinyon-juniper, ponderosa pine, at mid- to high-elevations, and salt desert shrub at low elevations. Low sagebrush or black sagebrush may form large islands within this community where soils are shallow or have restrictive layers.

Concerns: Post-settlement conversion to cheatgrass is common and results in change in fire frequency and vegetation dynamics. Fire suppression can lead to pinyon-juniper encroachment with subsequent loss of shrub and herbaceous understory. Disturbance of this community may result in establishment of annual grasslands (e.g., cheatgrass) and/or noxious weeds. Lack of disturbance can result in pinyon-juniper encroachment where adjacent to pinyon-juniper woodlands.

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

Historic disturbance (fire) likely ranged from small (< 10 ac) to large (> 10,000 acres) depending on conditions, time since last ignition, and fuel loading. Assumed the average patch size is 250 acres.

Issues/Problems

1) Some reviewers recommended merging all Wyoming big sagebrush PNVGs: R2SBWY, R2SBWYse, and R2SBWYwt. These PNVGs do not occur in the same areas or effective precipitation zones. Revised PNVGs are more clearly distinguished with greater differences in MFIs and fire behavior. Also, some reviewers did not know the LANDFIRE definition of mixed severity fire (25-75% of vegetation within burn perimeter is top killed by fire), which caused them to include mixed severity within replacement fire (>75% topkill).

2) There are no data, although abundant opinions, for the percentage of replacement and mixed severity fires, especially during mid-development, or whether surface fires occurred at all during early development during the pre-settlement phase.

Model Evolution and Comments

This model assumes the sites are near pinyon-juniper woodlands and without frequent fire, the p-j will encroach into the sagebrush range site.

The first three development classes chosen for this PNVG correspond to the early, mid-, and late seral stages familiar to range ecologists. The two classes with conifer invasion (classes D and E) approximately correspond to Miller and Tausch's (2001) phases 2 and 3 of pinyon and juniper invasion into shrublands. A

PNVG for Wyoming big sagebrush without tree invasion (R2SBWy; due to low elevation) was developed.

Succession Classes

Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 15%

Early1 PostRep

Description

Post-replacement disturbance; grass dominated with scattered shrubs. Fuel loading discontinuous. Surface fire occurs every 200 years on average but has no effect on succession. Succession to class B after 20 years.

Indicator Species* and Canopy Position

ACHY
HECO8
CHVI8
ARTRW8

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	10 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 50%

Mid1 Open

Description

Shrubs and herbaceous can be co-dominant, fine fuels bridge the woody fuels, but fuel discontinuities are possible. Replacement fire accounts for 80% of fire activity (mean FRI of 125 years), whereas mixed severity fire occurs every 500 years on average (20% of fire activity) and maintains vegetation in class B. Succession to class C after 40 years.

Indicator Species* and Canopy Position

ARTRW8
ACHY
CHVI8
HECO26

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	11 %	25 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class C 25%

Mid2 Closed

Description

Shrubs dominate the landscape; fuel loading is primarily woody vegetation. Shrub density sufficient in old stands to carry the fire without fine fuels. Establishment of pinyon and juniper seedlings and saplings widely scattered. Replacement fire (mean FRI of 100 years) and rare flood events (return interval of 333 years) cause a transition to class A. Prolonged

Indicator Species* and Canopy Position

ARTRW8
CHVI8
ELEL5
HECO26

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	26 %	35 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

drought (mean return interval of 100 years) and insect/disease (every 75 years on average) cause a transition to class B. Succession to class D after 40 years.

Class D 5%

Late1 Open

Description

Pinyon-juniper encroachment where disturbance has not occurred for 100+ years (tree species cover <15%). Saplings and young trees are the dominant lifeform. Sagebrush cover (<25%) and herbaceous cover decreasing compared to class C. Replacement fire occurs every 125 years on average. Insect/disease (every 75 years) and prolonged drought (every 100 years) thin both trees and shrubs, causing a transition to class C. Succession to class E after 50 years.

Indicator Species* and Canopy Position

JUNIP
PIMO
ARTRW8

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	<i>Min</i>	<i>Max</i>
<i>Cover</i>	0 %	15 %
<i>Height</i>	no data	no data
<i>Tree Size Class</i>	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class E 5%

Late1 Closed

Description

Pinyon-juniper woodland (cover 16-90%) where disturbance does not occur for 50+ years in Class D. Shrub cover <10% and graminoids scattered. Replacement fire occurs every 125 years on average. Prolonged drought thins trees, causing a transition to class B. Succession from class E to E.

Indicator Species* and Canopy Position

JUNIP
PIMO

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	<i>Min</i>	<i>Max</i>
<i>Cover</i>	16 %	90 %
<i>Height</i>	no data	no data
<i>Tree Size Class</i>	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Disturbances

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Non-Fire Disturbances Modeled

- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other:

Fire Regime Group: 4

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

Historical Fire Size (acres)

Avg:
Min:
Max:

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
<i>Replacement</i>	137	30	200	0.0073	84
<i>Mixed</i>	1000			0.001	11
<i>Surface</i>	2500			0.0004	5
<i>All Fires</i>	115			0.0087	

References

Brown, J. K. and J. K. Smith, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Cronquist, A., A.H. Holmgren, N.H. Holmgren, J.L. Reveal, and P.K. Holmgren. 1994. Intermountain Flora: Vascular Plants of the Intermountain West, U.S.A. Asterales. Volume 5. New York Botanical Garden, Bronx, NY.

Gruell, G.E. 1999. Historical and modern roles of fire in pinyon-juniper. P. 24-28. In: S.B. Monsen & R. Stevens (compilers). Proceedings: ecology and management of pinyon-juniper communities within the Interior West; 1997, Provo, UT. Proc. RMRS-P-9. Ogden, UT. U.S. Dept. Ag., Forest Service, Rocky Mountain Research Station.

Kuchler, A.W. 1985. Potential natural vegetation (map at scale of 1:7,500,000). In: U.S. Geological survey, The National Atlas of the USA. U.S. Govt. Print. Off. Washington, D.C.

Miller, R.F. and J.A. Rose. 1999. Fire history and western juniper encroachment in sagebrush-steppe. Journal of Range Management. 550-559.

Miller, R.F. and L.L. Eddleman. 2000. Spatial and temporal changes of sage grouse habitat in the sagebrush biome. Oregon State Univ. Agr. Exp. Stat. Technical Bull. 151. 35pp.

Miller, R. F. and R. J. Tausch. 2001. The role of fire in juniper and pinyon woodlands: a descriptive analysis. Proceedings: The First National Congress on Fire, Ecology, Prevention, and Management. San Diego, CA, Nov. 27- Dec. 1, 2000. Tall Timbers Research Station, Tallahassee, FL. Miscellaneous Publication 11, p:15-30.

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

NRCS. 2003. Major Land Resource Area 28A Great Salt Lake Area. Nevada Ecological Site Descriptions. Reno, NV.

NRCS. 2003. Major Land Resource Area 28B Central Nevada Basin and Range. Nevada Ecological Site Descriptions. Reno, NV.

NRCS. 2003. Major Land Resource Area 25 Owyhee High Plateau. Nevada Ecological Site Descriptions. Reno, NV.

NRCS. 2003. Major Land Resource Area 24 Humboldt Area. Nevada Ecological Site Descriptions. Reno, NV.

NRCS. 2003. Major Land Resource Area 27 Fallon-Lovelock Area. Nevada Ecological Site Descriptions. Reno, NV.

Tausch, R.J. and R.S. Nowak. 1999. Fifty years of ecotone change between shrub and tree dominance in the Jack Springs Pinyon Research Natural Area. P.71-77. In: E.D. McArthur, W. K. Ostler, & C.L Wambolt (compilers). Proceedings: shrubland ecotones. 1998. Ephram, UT. Proc. RMRS-P-11. Ogden, UT. U.S. Dept. Ag., Forest Service, Rocky Mountain Research Station.

Tiltsdale, E.W. 1994. Great Basin region: sagebrush types. P. 40-46. In: T.N. Shiflet (ed.) Rangeland Cover Types. Soc. Range Manage., Denver, CO.

West, N.E. 1983. Western Intermountain sagebrush steppe. P. 351-297. In: N.E. West (ed.) Ecosystems of the World 5: Temperate deserts and semi-deserts. Elsevier Scientific Publishing Company, New York, NY.